

**MMWR***Weekly*

July 31, 2009 / 58(29);800-804

## Fatalities Caused by Cattle --- Four States, 2003--2008

During 2003--2007, deaths occurring in the production of crops and animals in the United States totaled 2,334; of these, 108 (5%) involved cattle as either the primary or secondary cause (1 ). During the same period, Iowa, Kansas, Missouri, and Nebraska accounted for 16% of the nation's approximately 985,000 cattle operations and 21% of the nation's cattle and calf herd (2 ). To better characterize cattle-caused deaths in these four states, investigators reviewed all such deaths occurring during the period 2003--2008 that were detected by two surveillance programs, the Iowa Fatality Assessment and Control Evaluation (IA FACE) and the Great Plains Center for Agricultural Health (GPCAH). This report summarizes that investigation, which identified 21 cattle-related deaths. These deaths occurred throughout the year, and decedents tended to be older (aged  $\geq 60$  years) (67%) and male (95%). Except in one case, the cause of death was blunt force trauma to the head or chest. Circumstances associated with these deaths included working with cattle in enclosed areas (33%), moving or herding cattle (24%), loading (14%), and feeding (14%). One third of the deaths were caused by animals that had previously exhibited aggressive behavior. To reduce the risk for death from cattle-caused injuries, farmers and ranchers should be aware of and follow recommended practices for safe livestock-handling facilities and proper precautions for working with cattle, especially cattle that have exhibited aggressiveness.

Data gathering and analysis were performed collaboratively by IA FACE (operated by the University of Iowa on behalf of the Iowa Department of Public Health) and GPCAH (part of the University of Iowa's College of Public Health). Both programs are funded by CDC and collect surveillance data on agricultural deaths.\* IA FACE collects basic information on all traumatic occupational fatalities in Iowa as identified primarily through multisource surveillance of the media, including newspapers, radio, television, and the Internet. Once alerted to a potential occupational death, IA FACE requests reports from investigating authorities such as the local police and sheriff's departments, emergency medical services, and the medical examiner. GPCAH surveillance is based solely on reports from Iowa, Kansas, Missouri, and Nebraska newspapers and other periodicals. Since 2003, GPCAH has been building a press report database, which includes descriptive information about the victim, event, circumstances, and nature of the injuries in fatal and nonfatal farm and agricultural injury events within the four states.

In this analysis, cases were defined as occupational fatalities caused by cattle that occurred in Iowa,

Kansas, Missouri, or Nebraska during 2003--2008. Fatalities that occurred when motor vehicles crashed into cattle on roadways (such as while cattle were being herded with an all-terrain vehicle or pickup truck in a pasture) were excluded.

## Surveillance Results

A total of 21 deaths met the case definition for 2003--2008 ([Table 1](#)). Four fatalities occurred in 2003, two in 2004, six in 2005, and three each year during 2006--2008. During these years, eight of the fatalities occurred in Iowa, two in Kansas, seven in Missouri, and four in Nebraska. The 21 decedents ranged in age from 8 to 86 years, with a median age of 65 years (mean age: 61 years) ([Table 2](#)). Only one of the victims was female. One of the victims was a boy aged 8 years who was helping castrate cattle when he was crushed against a squeeze chute. One third of the deaths occurred in March and April.

The victims' most common activities at the time of death were working with and treating cattle in enclosed spaces such as pens and chutes ( $n = 7$ ) and moving or sorting cattle toward pens, barns, or pastures ( $n = 5$ ). Incidents also occurred while loading cattle into trucks or trailers ( $n = 3$ ), feeding ( $n = 3$ ), or working in an open pasture ( $n = 3$ ).

Ten of the 21 fatalities involved attacks by individual bulls, six involved attacks by individual cows, and five involved multiple cattle. In seven attacks (whether witnessed or not), the bull or cow was known to have exhibited aggressive behavior in the past. In 16 of the cases, the animal was deemed to have purposefully struck the victim; five other deaths were caused by being crushed against a stationary object or struck by a gate (secondary to the action of cattle). All but one death resulted from blunt force trauma to the chest and/or head; one resulted from inadvertent injection of the antibiotic Micotil 300 (tilmicosin phosphate) from a syringe in the victim's pocket when he was knocked down by a cow.

## Illustrative Case Reports

The following case summaries illustrate the most common circumstances of the cases identified for this report.

**Case 1.** In August 2005, a woman in Missouri aged 65 years was removing a dead, newborn calf from a pasture when a cow knocked her down, stomped her, and butted her while she was lying on the ground. The coroner reportedly stated that death resulted from blunt force trauma to the woman's head and chest. No autopsy was performed.

**Case 2.** In November 2005, a man in Iowa aged 65 years was helping his son sort beef cattle for loading onto a truck. He was attempting to guide one of the animals toward the truck when it turned into him, crushing him against the barn door. According to witnesses, he stopped breathing immediately. The medical examiner's report stated that death was caused by blunt force trauma to the man's chest.

**Case 3.** In April 2006, a man in Iowa aged 63 years was herding cattle into his dairy barn for milking when a

bull came into the barn and repeatedly butted him, pinned him against a fence, and stomped him. According to the attending physician's death record, the man sustained multiple rib fractures, lacerated pulmonary arteries, and head injuries. The man's family said that the bull was known to be dangerous and had been threatening in the past.

**Case 4.** In August 2007, a man in Iowa aged 45 years who was working alone in a pasture was attacked by a bull that had been bottle-fed and raised by the family but, according to family members, had become more aggressive recently. The attack was not witnessed, but the man was able to call his wife for assistance on his cell phone before he died and told her he had been attacked. According to the state medical examiner's autopsy report, he died of blunt force injuries to the chest.

**Reported by:** *WT Sanderson, PhD,  
MD Madsen, MBA, Great  
Plains Center  
for Agricultural Health  
and the Injury  
Prevention Research  
Center, College of  
Public Health, Dept  
of Occupational  
and Environmental  
Health, Univ of Iowa,  
Iowa City.*

### **Editorial Note:**

Large livestock are powerful, quick, protective of their territory and offspring, and especially unpredictable during breeding and birthing periods (3--5 ). Mothering livestock often protect their young aggressively. Dairy bulls, which have more frequent contact with humans than do beef cattle, are known to be especially possessive of their herd and occasionally disrupt daily feeding, cleaning, and milking routines (5 ). The findings in this report confirm earlier research substantiating the risk for death to farmers and ranchers from contact with cattle (3,5--8 ). Previously published reports have described the nature and frequency of cattle-related deaths and injuries. Among 739 patients admitted to a referral trauma center in Wisconsin during a 12-year period because of injuries incurred while farming, 30% involved injuries from farm animals (6 ). Working with bulls involves higher risk for injury. In a study of farm worker injuries based on surveillance data from New York, bulls were found to account for 25% of animal-related injuries (7 ). Among the deaths described in this report, four (19%) were caused by dairy bulls during feeding or milking operations.

Of the decedents mentioned in this report, 13 of 20 (65%) were men aged  $\geq 60$  years. The methodology used in this analysis did not allow the calculation of age-specific risks and could not determine whether this age and sex profile reflected the demographics of farmers involved in close contact with cattle in the four states, or

a greater risk for death among older farmers and ranchers. A case-control study of Iowa livestock farmers found that use of a hearing aid (odds ratio [OR] = 5.4) and doctor-diagnosed arthritis or rheumatism (OR = 3.0) were significantly associated with injuries related to animals (8). Age-related reduced hearing and reduced ability to react might contribute to this risk. Because approximately one third of the deaths described in this report occurred when the farmer was working alone, some of these deaths might have been prevented if a coworker had been present to help observe cattle behavior and movement and to provide prompt aid in case of injury. This might be especially useful when working with bulls or cows known to be aggressive, given that seven of the deaths described in this report involved such cattle.

The findings in this report are subject to at least two limitations. First, IA FACE surveillance, which involves more in-depth follow-up, only captured fatalities associated with work in Iowa. GPCAH surveillance, which is conducted in Iowa, Kansas, Missouri, and Nebraska, only captured accounts that appeared in newspapers or other periodicals. Therefore, reports from coroners or medical examiners, law enforcement, and emergency services were not obtained in Kansas, Missouri, or Nebraska. As a result, details about incidents in these three states often were limited (e.g., the age and sex of the decedent always were reported, but occasionally the decedent's activities and surroundings were not well reported). Second, reliance primarily on news reports means that some fatalities might go unreported. In Iowa, during 2003--2007, all seven of the fatalities caused by cattle that were documented by the state-based Census of Fatal Occupational Injuries (CFOI) of the U.S. Department of Labor's Bureau of Labor statistics also were captured through IA FACE and GPCAH surveillance. However, CFOI documented four cattle-caused fatalities in Kansas, seven in Missouri, and four in Nebraska, whereas GPCAH captured only two fatalities in Kansas, six in Missouri, and two in Nebraska. These data indicate that in states where only press clips were used to document agricultural fatalities, five out of 15 (33%) of the fatalities were unreported, suggesting a sensitivity of 67%. However, the advantage of using press reports is that more information regarding the circumstances of the deaths might be collected. In published studies, the sensitivity of newspapers as an injury surveillance source has varied according to the type of injury (9).

Previously published reports have recommended that cattle handling facilities be designed for optimum safety, such as the placing of sturdy barriers between cattle and persons, allowing for directed movement of cattle, and providing means for rapid exit from the cattle area (10). Information on safe cattle handling and safe cattle-handling facilities is available from the National Agricultural Safety Database at <http://www.nasdonline.org/menu/topic/animals.html>.

## Acknowledgments

This report is based, in part, on contributions by participating state agencies; Bureau of Labor Statistics staff; FACE program staff; J Kraemer, Iowa Office of the State Medical Examiner; K Leinenkugel, Occupational Safety and Health Surveillance Program, Iowa Dept of Public Health; and JR Myers, National Institute for Occupational Safety and Health, CDC.

## References

1. US Department of Labor, Bureau of Labor Statistics. Census of Fatal Occupational Injuries---current and revised data, 2003--2007. Available at <http://www.bls.gov/iif/oshcfoi1.htm>.
2. U.S. & all states data: cattle & calves, cattle operations, 2003--2007. National Agricultural Statistics Database. Available at [http://www.nass.usda.gov/QuickStats/PullData\\_US.jsp](http://www.nass.usda.gov/QuickStats/PullData_US.jsp).
3. Dogan KH, Serafettin D, Erkol E, Sunam G, Kucukkartallar T. Injuries and deaths occurring as a result of a bull attack. J Agromedicine 2008;13:191--6.
4. Grandin T (ed). Livestock handling and transport. 3rd edition. Wallingford, United Kingdom: Oxford University Press; 2007.
5. Boyle D, Gerberich S, Gibson R, et al. Injury from dairy cattle activities. Epidemiology 1997;8:37--41.
6. Cogbill T, Steenlage E, Landercasper J, Strutt P. Death and disability from agricultural injuries in Wisconsin: a 12-year experience with 739 patients. J Trauma 1991;31:1632--7.
7. Casey G, Grant A, Roerig D, et al. Farm worker injuries associated with bulls. New York State 1991--1996. AAOHN J 1997;45:393--6.
8. Sprince N, Park H, Zwerling C, et al. Risk factors for animal-related injury among Iowa large-livestock farmers: a case-control study nested in the agricultural health study. J Rural Health 2003;19:165--73.
9. Rainey DY, Runyan CW. Newspapers: a source for injury surveillance? Am J Public Health 1992;82:745--6.
10. Bicudo JR, McNeill S, Turner L, Burris R, Anderson J. Cattle handling facilities: planning, components, and layouts. Lexington, KY: Kentucky Cooperative Extension Service, University of Kentucky College of Agriculture; 2002. Available at <http://www.ca.uky.edu/agc/pubs/aen/aen82/aen82.pdf>.

\* Additional information about IA FACE is available at <http://www.public-health.uiowa.edu/face>. Information on GPCAH is available at <http://www.public-health.uiowa.edu/gpcah>.

**TABLE 1. Characteristics of cattle-caused fatalities --- Iowa, Kansas, Missouri, and Nebraska, 2003--2008\*†**

Month and year	State	Decedent	Sex	Age (yrs)	Animal involved	Incident
Mar 2003	IA	Cattle farmer	Male	77	Beef cattle	Struck by gate when cattle charged while being herded
Oct 2004	IA	Cattle farmer	Male	48	Beef cattle	Pinned against barn wall while working with cattle
Nov 2004	IA	Dairy farmer	Male	77	Dairy bull	Attacked from behind by bull when feeding dairy cows
Sep 2005	IA	Veterinarian	Male	64	Beef bull	Attacked by bull when vaccinating and applying insecticide on cattle
Nov 2005	IA	Cattle farmer	Male	65	Beef cattle	Crushed against barn door when sorting cattle
Apr 2006	IA	Dairy farmer	Male	65	Dairy bull	Attacked by bull when herding cows for milking
Apr 2006	IA	Dairy farmer	Male	63	Dairy bull	Attacked by bull while moving cows into milking parlor
Aug 2007	IA	Cattle farmer	Male	45	Beef bull	Attacked by bull when alone in pasture
Apr 2003	KS	Cattle farmer	Male	86	Beef calves	Knocked steel gate on top of himself while loading calves onto a trailer
Jul 2005	KS	Cattle farmer	Male	74	Beef bull	Trampled by bull being moved from one pasture to another

Mar 2003	MO	Cattle farmer	Male	71	Beef cows	Found fatally injured in pen with two cows and newborn calf
Feb 2005	MO	Cattle farmer	Male	62	Beef cow	Kicked in head by cow
Aug 2005	MO	Cattle farmer	Female	65	Beef cow	Attacked by cow when removing dead calf from pasture
Dec 2005	MO	Cattle farmer	Male	53	Beef bull	Mauled by aggressive bull in pasture while retrieving cows
Jan 2006	MO	Dairy farmer	Male	39	Dairy bull	Mauled and crushed against barn wall by bull while feeding cows
Sep 2007	MO	Cattle farmer	Male	75	Beef bull	Gored while loading bull into trailer
Jan 2008	MO	Cattle farmer	Male	72	Beef bull	Rammed by bull while feeding cattle
Mar 2003	NE	Cattle farmer	Male	38	Beef cow	Injected with Micotil from syringe in his pocket when cow pushed him down
Mar 2007	NE	Cattle farmer	Male	47	Beef cow	Crushed in pen when attacked by cow with calf
May 2008	NE	Cattle farmer	Male	81	Beef cow	Attacked by cow while working in pen
Jun 2008	NE	Child <sup>§</sup>	Male	8	Beef cattle	Crushed while moving cattle through squeeze chute

\* Based on cases identified through the Iowa Fatality Assessment and Control Evaluation (IA FACE) (operated by the University of Iowa on behalf of the Iowa Department of Public Health) and the Great Plains Center for Agricultural Health (GPCAH) (part of the University of Iowa's College of Public Health). IA FACE collects basic information on all traumatic occupational fatalities in Iowa as identified primarily through multisource surveillance (by IA FACE staff and professional colleagues across the state) of the media, including newspapers, radio, television, and the internet. Once alerted to a potential occupational death, IA FACE requests reports from investigating authorities such as the local police and sheriff's departments, emergency medical services, and medical examiner. GPCAH surveillance is based solely on reports from Iowa, Kansas, Missouri, and Nebraska newspapers and other periodicals. Additional information about IA FACE is available at <http://www.public-health.uiowa.edu/face>. Information on GPCAH is available at <http://www.public-health.uiowa.edu/gpcah>.

† Cases were defined as occupational fatalities caused by cattle that occurred in Iowa, Kansas, Missouri, or Nebraska during 2003--2008. Fatalities that occurred when motor vehicles crashed into cattle on roadways (such as while cattle were being herded with an all-terrain vehicle or pickup truck in a pasture) were excluded.

§ Child was killed while helping on the family farm.

**TABLE 2. Number and percentage of cattle-caused fatalities, by selected characteristics --- Iowa, Kansas, Missouri, and Nebraska, 2003--2008\*†**

Characteristic	No.	(%) <sup>§</sup>
<b>Sex of decedent</b>		
Male	20	(95)
Female	1	(5)
<b>Age group (yrs) of decedent</b>		
<60	7	(33)
≥60	14	(67)

<b>Operation/Activity</b>		
Herding/Moving/Sorting	5	(24)
Loading	3	(14)
Feeding	3	(14)
Tending/Treating in enclosed area	7	(33)
Attacked in open pasture	3	(14)
<b>Animal involved</b>		
Bull	10	(48)
Cow with calf	3	(14)
Cow (no calf)	3	(14)
Multiple cattle	5	(24)
<b>Total</b>	<b>21</b>	<b>(100)</b>

\* Based on cases identified through the Iowa Fatality Assessment and Control Evaluation (IA FACE) (operated by the University of Iowa on behalf of the Iowa Department of Public Health) and the Great Plains Center for Agricultural Health (GPCAH) (part of the University of Iowa's College of Public Health). IA FACE collects basic information on all traumatic occupational fatalities in Iowa as identified primarily through multisource surveillance (by IA FACE staff and professional colleagues across the state) of the media, including newspapers, radio, television, and the internet. Once alerted to a potential occupational death, IA FACE requests reports from investigating authorities such as the local police and sheriff's departments, emergency medical services, and medical examiner. GPCAH surveillance is based solely on reports from Iowa, Kansas, Missouri, and Nebraska newspapers and other periodicals. Additional information about IA FACE is available at <http://www.public-health.uiowa.edu/face>. Information on GPCAH is available at <http://www.public-health.uiowa.edu/gpcah>.

† Cases were defined as occupational fatalities caused by cattle that occurred in Iowa, Kansas, Missouri, or Nebraska during 2003--2008. Fatalities that occurred when motor vehicles crashed into cattle on roadways (such as while cattle were being herded with an all-terrain vehicle or pickup truck in a pasture) were excluded.

§ Percentages might not sum to 100% because of rounding.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites. URL addresses listed in *MMWR* were current as of the date of publication.

All *MMWR* HTML versions of articles are electronic conversions from typeset documents. This conversion might result in character translation or format errors in the HTML version. Users are referred to the electronic PDF version (<http://www.cdc.gov/mmwr>) and/or the original *MMWR* paper copy for printable versions of official text, figures, and tables. An original paper copy of this issue can be obtained from the Superintendent of Documents, U.S. Government Printing Office (GPO),

Washington, DC 20402-9371; telephone: (202) 512-1800. Contact GPO for current prices.

\*\*Questions or messages regarding errors in formatting should be addressed to [mmwrq@cdc.gov](mailto:mmwrq@cdc.gov).

Date last reviewed: 7/30/2009

---

[HOME](#) | [ABOUT MMWR](#) | [MMWR SEARCH](#) | [DOWNLOADS](#) | [RSS](#) | [CONTACT](#)  
[POLICY](#) | [DISCLAIMER](#) | [ACCESSIBILITY](#)

**SAFER • HEALTHIER • PEOPLE™**

**Morbidity and Mortality Weekly Report**  
Centers for Disease Control and Prevention  
1600 Clifton Rd, MailStop E-90, Atlanta, GA 30333, U.S.A



Department of Health  
and Human Services